

Extent of disease affects the usefulness of fecal biomarkers in ulcerative colitis

Authors:

Akihito Sakuraba¹, Nobuki Nemoto¹, Noritaka Hibi¹, Ryo Ozaki¹, Sotaro Tokunaga¹, Oki Kikuchi¹, Shintaro Minowa¹, Tatsuya Mitsui¹, Miki Miura¹, Daisuke Saito¹, Mari Hayashida¹, Jun Miyoshi¹, Minoru Matsuura¹, Masayoshi Yoneyama², Hiroaki Ohnishi³, Tadakazu Hisamatsu^{1*}

Affiliation:

- 1) Department of Gastroenterology and Hepatology, Kyorin University School of Medicine, Tokyo, Japan
- 2) Department of Clinical Laboratory, Kyorin University Hospital, Tokyo, Japan.
- 3) Department of Laboratory Medicine, Kyorin University School of Medicine, Tokyo, Japan.

*** Corresponding author:**

Tadakazu Hisamatsu, M.D., Ph.D., FACG, AGAF
Department of Gastroenterology and Hepatology,
Kyorin University School of Medicine.
6-20-2 Shinkawa, Mitaka-shi, Tokyo 181-8611, Japan

Tel.: +81-422-47-5511 (ext. 5279)

Fax: +81-422-44-0655

E-mail: [thisamatsu@ks.kyorin-u.ac.jp](mailto>thisamatsu@ks.kyorin-u.ac.jp)

Running title: Disease extent and fecal biomarkers in UC

Supplemental Figure 2


a

All cases


All cases

All cases


Fecal calprotectin (mg/kg)



Fecal immunochemical test (ng/ml)



Fecal lactoferrin (mg/kg)




b

Pancolitis


Pancolitis

Pancolitis


Fecal calprotectin (mg/kg)



Fecal immunochemical test (ng/ml)



Fecal lactoferrin (mg/kg)




c

Left-sided colitis


Left-sided colitis

Left-sided colitis


Fecal calprotectin (mg/kg)



Fecal immunochemical test (ng/ml)



Fecal lactoferrin (mg/kg)




d

Proctitis


Proctitis

Proctitis

Fecal calprotectin (mg/kg)



Fecal immunochemical test (ng/ml)



Fecal lactoferrin (mg/kg)

